


Lecture 7 Quasi-Geostrophic (QG) Approximation and QG Vorticity Equation

7.1 Quasi-Geostrophic Approximation

(Equation editor: $D/Dt = \partial/\partial t + u\partial/\partial x$)

Purpose: The primitive equations contain several terms of secondary significance, thus can be simplified by making the **quasi-geostrophic (QG) approximation** to help understand the basic dynamics of synoptic atmospheric motions in midlatitudes.



QG approximation will lead to two major equations:

- Geopotential height tendency equation for predicting the height tendency
- Omega equation for diagnose the vertical motion

➤ QG Approximation of the momentum equations

Consider the primitive equations in isobaric coordinates

$$\frac{Du}{Dt} = fv - \frac{\partial \phi}{\partial x} \quad x\text{-momentum equation}$$

$$\frac{Dv}{Dt} = -fu - \frac{\partial \phi}{\partial y} \quad y\text{-momentum equation}$$

$$\frac{\partial \phi}{\partial p} = -\frac{RT}{p} \quad \text{Hydrostatic equation}$$

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial \omega}{\partial p} = 0 \quad \text{Continuity equation}$$

$$\frac{DT}{Dt} = \left(\frac{RT}{c_p p} \right) \omega + \left(\frac{1}{c_p} \right) \frac{DQ}{Dt} \quad \text{Thermodynamic equation}$$

➤ We have already made the following assumptions or approximations:

- (1) Inviscid
- (2) Hydrostatic
- (3) No Earth curvature effect

➤ Next, the total horizontal velocity is decomposed into geostrophic and ageostrophic components

$$u = u_g + u_a \quad \text{and} \quad v = v_g + v_a, \text{ or in vector form}$$

$$\mathbf{V} = \mathbf{V}_g + \mathbf{V}_a$$

(4) Assume $u_a \ll u_g$ and $v_a \ll v_g$, which gives

$$\frac{Du}{Dt} = \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + \omega \frac{\partial u}{\partial p} \approx \frac{D_g u_g}{Dt},$$

$$\frac{Dv}{Dt} = \frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + \omega \frac{\partial v}{\partial p} \approx \frac{D_g v_g}{Dt},$$

where $\frac{D_g}{Dt} = \frac{\partial}{\partial t} + u_g \frac{\partial}{\partial x} + v_g \frac{\partial}{\partial y}$.

It is valid only when $R_o \ll 1$.

(5) Make midlatitude β -plane approximation: $f = f_o + \beta y$ where β is a constant.

(6) Assume $f = f_o$ in the geostrophic wind balance

$$u_g \equiv -\frac{1}{f_o} \frac{\partial \phi}{\partial y}, \quad v_g \equiv \frac{1}{f_o} \frac{\partial \phi}{\partial x}$$

With the above assumptions and approximations, the x -momentum equation becomes

$$\frac{\partial u_g}{\partial t} + u_g \frac{\partial u_g}{\partial x} + v_g \frac{\partial u_g}{\partial y} = -\frac{\partial \phi}{\partial x} + (f_o + \beta y)(v_g + v_a) \approx f_o v_a + \beta y v_g$$

$$\frac{\partial v_g}{\partial t} + u_g \frac{\partial v_g}{\partial x} + v_g \frac{\partial v_g}{\partial y} = -\frac{\partial \phi}{\partial y} - (f_o + \beta y)(u_g + u_a) \approx -f_o u_a - \beta y u_g$$

In summary, the QG approximation neglects the following effects:

- Friction
- Horizontal advection of momentum by the ageostrophic wind, e.g. $\left(u_a \frac{\partial u_g}{\partial x}, v_a \frac{\partial u_g}{\partial y}\right)$, & on ageostrophic wind, e.g. $\left(u_a \frac{\partial u_a}{\partial x}, v_a \frac{\partial u_a}{\partial y}\right)$
- Vertical advection of momentum, e.g. $\left(\omega \frac{\partial u_g}{\partial p}, \omega \frac{\partial u_a}{\partial p}\right)$
- Local changes in the ageostrophic wind, e.g. $\frac{\partial u_a}{\partial t}$
- Advection of the ageostrophic momentum by the geostrophic wind, e.g. $\left(u_g \frac{\partial u_a}{\partial x}, v_g \frac{\partial u_a}{\partial y}\right)$

➤ QG Continuity Equation

Substituting $u = u_g + u_a$ and $v = v_g + v_a$ into

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial \omega}{\partial p} = 0$$

and using

$$u_g \equiv -\frac{1}{f_o} \frac{\partial \phi}{\partial y}, \quad v_g \equiv \frac{1}{f_o} \frac{\partial \phi}{\partial x}, \quad \text{and} \quad \frac{\partial u_g}{\partial x} + \frac{\partial v_g}{\partial y} = 0 \quad \text{leads to}$$

$$\frac{\partial u_a}{\partial x} + \frac{\partial v_a}{\partial y} + \frac{\partial \omega}{\partial p} = 0$$

➤ QG Thermodynamic Equation

The thermodynamic energy equation in the primitive set of equations

$$\frac{DT}{Dt} = \left(\frac{RT}{c_p p} \right) \omega + \left(\frac{1}{c_p} \right) \frac{DQ}{Dt}$$

can be rewritten as

$$\frac{\partial T}{\partial t} + u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} - \left(\frac{\sigma p}{R} \right) \omega = \frac{1}{c_p} \frac{DQ}{Dt} \quad \text{where} \quad \sigma = - \frac{RT}{\theta p} \frac{\partial \theta}{\partial p}.$$

Applying the primary QG approximation ($u \approx u_g$ and $v \approx v_g$) leads to

$$\frac{\partial T}{\partial t} = -u_g \frac{\partial T}{\partial x} - v_g \frac{\partial T}{\partial y} + \left(\frac{\sigma p}{R} \right) \omega + \frac{J}{c_p}$$

In summary, the QG equations can be written as

$$\frac{\partial u_g}{\partial t} + u_g \frac{\partial u_g}{\partial x} + v_g \frac{\partial u_g}{\partial y} = f_0 v_a + \beta y v_g \quad (6.10)$$

$$\frac{\partial v_g}{\partial t} + u_g \frac{\partial v_g}{\partial x} + v_g \frac{\partial v_g}{\partial y} = -f_0 u_a - \beta y u_g \quad (6.11)$$

$$\frac{\partial \phi}{\partial p} = - \frac{RT}{p} \quad (6.2)$$

$$\frac{\partial u_a}{\partial x} + \frac{\partial v_a}{\partial y} + \frac{\partial \omega}{\partial p} = 0 \quad (6.12)$$

$$\frac{\partial T}{\partial t} = -u_g \frac{\partial T}{\partial x} - v_g \frac{\partial T}{\partial y} + \left(\frac{\sigma p}{R} \right) \omega + \frac{J}{c_p} \quad (6.13)$$

$$u_g = - \frac{1}{f_0} \frac{\partial \phi}{\partial y} \quad (3/16/17)$$

$$v_g = \frac{1}{f_0} \frac{\partial \phi}{\partial x} \quad (6.7)$$

7.2 Quasi-Geostrophic Vorticity Equation

➤ Start with QG equations of motion

$$\frac{\partial u_g}{\partial t} + u_g \frac{\partial u_g}{\partial x} + v_g \frac{\partial u_g}{\partial y} = f_0 v_a + \beta y v_g \quad (6.10)$$

$$\frac{\partial v_g}{\partial t} + u_g \frac{\partial v_g}{\partial x} + v_g \frac{\partial v_g}{\partial y} = -f_0 u_a - \beta y u_g \quad (6.11)$$

Taking cross differentiation of the above equations leads to the QG vorticity equation

$$\frac{\partial \zeta_g}{\partial t} + u_g \frac{\partial \zeta_g}{\partial x} + v_g \frac{\partial \zeta_g}{\partial y} = f_0 \frac{\partial \omega}{\partial p} - \beta v_g \quad (6.18)$$

where

$$\zeta_g = \frac{\partial v_g}{\partial x} - \frac{\partial u_g}{\partial y} = \frac{1}{f_0} \nabla^2 \phi \quad (6.15)$$

Equation (6.18) can be rewritten as

$$\frac{\partial \zeta_g}{\partial t} = -V_g \cdot \nabla (\zeta_g + f) + f_0 \frac{\partial \omega}{\partial p} \quad (6.18)'$$

Physical meaning of (6.18)':

LHS: local rate of change of geostrophic relative vorticity

RHS: (1) advection of absolute vorticity by geostrophic wind

(2) vorticity stretching by planetary vorticity

Substituting (6.12) into the equation leads to an alternative form of the **QG vorticity equation**

$$\frac{\partial \zeta_g}{\partial t} = -u_g \frac{\partial \zeta_g}{\partial x} - v_g \frac{\partial \zeta_g}{\partial y} - f_0 \left(\frac{\partial u_a}{\partial x} + \frac{\partial v_a}{\partial y} \right) - \beta v_g \quad (6.18)$$

Thus, ζ_g can be predicted if ϕ and the right-hand side of is known.