


Lecture 12 The 2nd Law of Thermodynamics

(Sec.3.6 of Hess – 2nd Law of Thermodynamics)

[For classical equation editor: ($dq=0$)]

➤ **Claim:** In a Carnot cycle, $q_2 / q_1 = T_2 / T_1$.



Proof:

Applying the first law of thermodynamics ($dq=c_vdT+pd\alpha$) to the isothermal processes $a \Rightarrow b$ and $c \Rightarrow d$ leads to

$$q_1 = \int_a^b p d\alpha = \int_a^b \frac{RT_1}{\alpha} d\alpha = RT_1 \ln \frac{\alpha_b}{\alpha_a} \quad (12.1)$$

$$q_2 = - \int_c^d p d\alpha = - \int_c^d \frac{RT_2}{\alpha} d\alpha = RT_2 \ln \frac{\alpha_c}{\alpha_d} \quad (12.2)$$

Equations (12.1) and (12.2) gives

$$\frac{q_2}{q_1} = \frac{RT_2 \ln(\alpha_c/\alpha_d)}{RT_1 \ln(\alpha_b/\alpha_a)} = \left(\frac{T_2}{T_1} \right) \frac{\ln(\alpha_c/\alpha_d)}{\ln(\alpha_b/\alpha_a)}.$$

Therefore, in order to have $q_2 / q_1 = T_2 / T_1$, it requires

$$\frac{\alpha_c}{\alpha_d} = \frac{\alpha_b}{\alpha_a}. \quad (12.3)$$

Applying the equation of state,

$$p\alpha = RT$$

to the initial and final states of the isothermal processes $a \Rightarrow b$ and $d \Rightarrow a$ and the Poisson's equation,

$$p\alpha^\gamma = \text{constant}$$

to the adiabatic processes $b \Rightarrow c$ and $d \Rightarrow a$ gives

$$p_a\alpha_a = p_b\alpha_b = RT_1 \quad (1)$$

$$p_b\alpha_b^\gamma = p_c\alpha_c^\gamma \quad (2)$$

$$p_c\alpha_c = p_d\alpha_d = RT_2 \quad (3)$$

$$p_d\alpha_d^\gamma = p_a\alpha_a^\gamma. \quad (4)$$

Eq. (3) gives

$$\frac{\alpha_c}{\alpha_d} = \frac{p_d}{p_c} \quad (5)$$

Eqs. (2) and (4) gives

$$P_c = P_b(\alpha_b^\gamma/\alpha_c^\gamma)$$

$$P_d = P_a(\alpha_a^\gamma/\alpha_d^\gamma)$$

Substituting p_c and p_d into (5) yields

$$\frac{\alpha_c}{\alpha_d} = \left(\frac{p_a}{p_b}\right) \left(\frac{\alpha_a^\gamma\alpha_c^\gamma}{\alpha_b^\gamma\alpha_d^\gamma}\right). \quad (7)$$

Eq. (1) gives

$$p_a/p_b = \alpha_b/\alpha_a \quad (7)$$

Substituting (7) into (6) leads to

$$\frac{\alpha_c}{\alpha_d} = \frac{\alpha_b}{\alpha_a}, \quad (12.3)$$

which proves $q_2/q_1 = T_2/T_1$.

➤ *The Second Law of Thermodynamics*

The above claim leads to the 2nd Law of Thermodynamics, which concerns about the maximum fraction of a quantity of heat that can be converted into useful work.

From the expression of the efficiency of heat engine:

$$\eta = 1 - \frac{q_2}{q_1} = 1 - \frac{T_2}{T_1}$$

Therefore, in order to have a 100% efficiency ($\eta = 1$), T_2 must be zero ($T_2 = 0 \text{ K}$). Is it possible? The answer is “no”. This leads to the Second Law of Thermodynamics: “it is impossible to make an engine that has a 100% efficiency.”

The 2nd law of thermodynamics (Kelvin-Planck statement):

It is impossible to construct an engine which operates in a cycle and which produces no other effect than the extraction of heat from a heat reservoir and the performance of an equivalent amount of work.

Alternative statement:

Heat cannot of itself (i.e., without the performance of work by some external agency) pass from a cold to a warm body.